

THE EFFECT OF HONEY BEE COLONY EFFICIENCY UPON CROP YIELD BESIDES FARMER BENEFIT

Dr. Ansuman Chatterjee¹, Anitha Nallasivam², Roopashree Rangaswamy³

^{1,2,3} Department of Agriculture, JAIN (Deemed-to-be University), Bengaluru, India

Email Id- ¹dr.ansumanchatterjee@cms.ac.in, ²dr.anithanallasivam@cms.ac.in, ³r.roopashree@jainuniversity.ac.in

Abstract

Despite widespread interest in pollinators' position in nutriment manufacture, their effect upon agriculturalists' benefit, that decides their subsistence as well as land-utilization choices, remains unknown. While normal pollinator assistances were commonly presumed, even within the same region, here was the latent for significant longitudinal distinction amongst yield types besides variations, as well as amongst pollinator administration policies. It investigated efficiency of honey bee colony utilized for pollination service, that includes pathogen control and non-natural winter feeding, affects flowers visitation, farmer benefit, and fruit production (Patagonia). Flower visiting rates were 130.5 percent higher in high-quality apple colonies than in traditional colonies resulting in a 70.5% rise in farm income. Colony consistency only influenced fruit weight, varieties a rise in farmer income to the same degree as the apple. It found no, in comparison to other reports, demonstrating the vulnerability mechanism. Our research found that simple improvements leased honey bee colony would increase visitation rates, increasing and thus increasing agriculturalists' incomes outpacing availability of honey bee colonies, our findings indicate practises intended to enhancing cluster efficiency could be able to help address this possible colony number shortfall. *A. mellifera* is the sole pollinator. With the reported reduction of bees in world, condition like this could sadly spread..

Key words: *Apis mellifera*, Apples, Colony quality, Farmer profits, Generalized multilevel path model, Pears, Pollination.

Introduction

Pollination as an ecology utility can be jeopardised by continuing pollinator decline. In several parts of the world, wild bee populations, which are important for crop pollination, have been declining. Despite the fact that the *Apis mellifera* has grown in grown faster rate divergent patterns can in pollination service. Because sufficient, the entomophiles output can eventually stop, or even transform into costs. As a result, there is a pressing create improving productivity while preserving biodiversity. Despite widespread interest in pollinators' position in nutriment manufacture, their effect upon agriculturalists' benefit, that decides their subsistence as well as land-utilization choices, remains unknown. While normal pollinator assistances were commonly presumed, even within the same region, here was the latent for significant longitudinal distinction amongst yield types besides variations, as well as amongst pollinator administration policies.

It investigated efficiency of honey bee colony utilized for pollination service, that includes pathogen control and non-natural winter feeding, affects flowers visitation, farmer benefit, and fruit production (Patagonia). Flower visiting rates were 130.5 percent higher in high-quality apple colonies than in traditional colonies resulting in a 70.5% rise in farm income. Colony consistency only influenced fruit weight, varieties a rise in farmer income to the same degree as the apple. It found no, in comparison to other reports, demonstrating the vulnerability mechanism. Our research found that simple improvements leased honey bee colony would increase visitation rates, increasing and thus increasing agriculturalists' incomes outpacing availability of honey bee colonies, our findings indicate practises intended to enhancing cluster efficiency could be able to help address this possible colony number shortfall. *A. mellifera* is the sole pollinator. With the reported reduction of bees in world, condition like this could sadly spread.

The feasibility of existing pollination procedures must be tested as a first step can be influenced by improving pollination through successful management. Pollination is important not only for total yield, but also for primary

controlled, little is known about honey bees' contribution to farmer incomes. Understanding the importance of essential for developing management strategies that improve pollination while reducing production variability and farmer earnings. However, to determine impacts of pollination service, it is essential to relate pollination activities to a farmer's benefit. The majority of research to date have concentrated on the impact of various pollinator control systems on yield quantity and efficiency, with only a handful addressing the economic implications.

Many of these experiments are limited in their ability to change processes in various applicable contexts due to their lack of an economic component [1]. Some most commercially important crops in some countries are apples (*Malus domestic*) and pears (*Pyrus communis*), which became the most exported fruits for many years. Numerous uninhabited bud tourists, such as solitary bees and bumble bee, are known to be effective pollinators of pears and apples. Several controlled pollinator insects, mainly on a regular basis [2]. In some other countries, it investigated the impact of honey bee cluster control, specifically colony planning as well as wellbeing, upon apple and pear pollination. It created a method to evaluate the effects of honey bee cluster administration on fruit quality and quantity, as well as relate to farmers' incomes. It show that honey bee cluster administration is especially important in agricultural ecosystem, especially when different pollinator like uninhabited bee is unavailable [3].

C. Brittain *et al.* performed this research the Valle area produces 74 and 86 percent of Argentina's pear and apple crops, respectively. It chose a 4.9 km and 29.5 km long wide field of 24–44 ha woods of mixed pear as well as apple cultivation traditional centred at approx. 37°38' S, 67°59' O. Herbicides, fungicides, and insecticides were used extensively in orchard management in the past added to apple vegetations to induce the abortion of misshapen apples. Since pear trees naturally contain thinning hormones, this procedure was not used on them. Farms had similar wood administration methods [4]. It chose an entire of 52 pears as well as 36 apple trees in the sample field, spaced at least 199 metres apart and spread over 89 planted plots of equal sizes (1.25 ha) nested within 21 dissimilar farms.

It chose the Red Delicious apple variety (38 trees) and the Abate Fetel (24 tree) and Packham Triumph (27 tree) pear variety as our focal trees because they are the greatest illustrative in this fruit-growing area. Self-incompatibility exists between Red Delicious as well as Packham Triumph, and Abate Fetel is partly self-fertile (4–11 percent). The Abate Fetel variety bloomed in September, during the flowering season. As a surrogate for cross-pollination potential [5]. At the start of the blossoming season for fruit tree in the study field, honey bee colonies are normally added to the orchards. Honey bee colonies are introduced in are distributed per plot by farmers. The average recommended colony density in our research area was between 4 and 8 apple and pear trees both have a ha1 value. In 9 of the 23 sample orchards, it implemented, and in the other 11 groves, it left the agriculturalists to handle fertilization utilizing traditional colony.

Unlike traditional colony, high-quality colony is made according to a set of guidelines. First, queens were induced to begin laying eggs earlier by feeding sugar syrup to colonies immediately subsequently the winter. Next, each colony's wellbeing is closely tracked after it was delivered. The colony is independent of African as well as Asian foulbrood, had a Varroa destructor infestation rate of less than 6% then are handled as needed to keep their wellbeing. As a result, when these colonies were released into the wild, they had a resting empress through a colony of minimum 19,999 bees [6]. It surveyed traditional as well as superior colony on one occasion in seven days throughout the blossoming season of pear as well as apple tree (in total 1000 colonies were surveyed). It counted the number of frames covered in bees at each survey to estimate colony strength.

Conventional colonies had half the number of frames filled by bees as superior colony ($F = 132$; $P = .002$, mean $sd = 3.9$, 1 vs 8.9, 0.9 for conservative as well as superior colony, respectively). The worth a agriculturalist had to wage for colonies rent (4.9 US\$ for a traditional colonies as well as 20.1 US\$ for a superior colonies) represented these disparities [7]. Since the behaviour of *A. mellifera* in cultured grounds decreases dramatically after a few hundred metres, this distance was selected (Fig. 1). A plot with a radius of 201 metres would cover the majority of the foraging honey bees that may visit a given focal tree. In addition, it calculated the linear distance (m) between each focal tree and the nearest colony [8].

It performed bee censuses at each of the 87 focal trees during the flowering time of each variety. Depending on logistics and environmental conditions, it performed 11-minute reflection cycles on all trees during its blossoming season, totalling 258 11-minute census upon 87 focal plants. Starting of all census tree1 and estimated how many flowers each floral visitor saw over the course of the next few weeks flowers with both had tree1. As a result, it was able to calculate it couldn't tell the difference between feral colonies and honey bees from controlled, so all *A. mellifera* visit were combined [9]. There is a difference in colony size. the median is represented by the dark line in the case, while the mean is represented by the black point. the first and third characteristics are located at the bottom and top of each box, respectively.




Fig. 1: Among Traditional and High-Quality Colonies,

Its total amount of fruits was calculated tree1 by measuring the number of fruits on each of the five marked branch (the identical one utilised to determine the blossom tourist visitation frequency) as well as the number of branches and the mean are multiplied [9]. It harvested and analysed that were tracked throughout the blossoming season. It estimated the size of the fruit (circumference, maximum weight height and width), counted the quantity of carpels by spores (0.1–6), as well as calculated the honey content of the surface as estimates of fruit quality. A tape measure was used to calculate the circumference of the fruit at its widest point, a calliper was used to measure the width and height to the nearest 0.15 mm, a digital balance was used to measure the weight to the nearest 0.015 g, and a handheld refractometer was used to measure the sugar content [10].

Literature Survey

M. A. Aizen *et al.* tried to understand possibility of a global pollination epidemic threatening agricultural production has piqued the attention of scientists, policymakers, and the general public in recent months. To date, provincial or resident losses in pollinators or inadequate pollination for specific crop have been used as justification for a global crisis. The study of FAO data, on the other hand, shows that the worldwide populace of controlled honey-bee hives had risen by 44.9 percent over the former fifty years, implying that monetary globalisation, rather than biotic forces, is driving the growing demand aimed at farming pollination service. However, accessible information shows an abundant faster (>299 percent) rise in the proportion of farming which relies on animal pollination over the past fifty years, that is putting worldwide pollination capability

under strain. While the chief reason of commercial agriculture's increasing reliance on pollinators should be taken into account when developing agricultural and conservation policies [1].

T. D. Breeze *et al.* inspected pollinator declines and raised questions about the availability of pollination services to agriculture across Europe. At the same time, EU agricultural and biofuel initiatives have boosted the cultured zone of insect pollinated yields across the landmass. Utilizing information from 40 European nations, this learning shows that between 2005 and 2010, the suggested quantity of honeybees essential to deliver yield pollination in Europe increased 5 times faster than honeybee stocks. As a result, honeybee stocks in 21 countries surveyed were inadequate to meet >91 percent of demand. These findings raise questions about numerous nations' ability to manage by large-scale losses of uninhabited pollinators, as well as various crucial gap in existing knowledge of pollination facility demand and supply, highlighting the urgent need for further research into this subject [3].

S. A. Cunningham *et al.* explained there is mounting evidence that better pollination practises will help promote higher yields and lower inconsistency for a variety of insect pollinated yields around the world. Succeeded honeybee is used to pollinate certain wood yields, but it is less widely utilized to pollinate ground yields, partly due to a lack of certainty about whether the possible gains outweigh the costs. Since there aren't enough experiments done at the right size, there's a lot of confusion. The yield benefits of controlled honeybees added to *Vicia faba* in field scale trials are investigated in this report. We placed honeybee hives in 16 fields in South Australia and found that as the distance between the hives increased, so did bee behaviour and the number of fruits per stem. Utilizing crop map information obtained at harvest, we investigated the spatial pattern of yield and discovered that distance from hives had an impact on mean crop (which decreased) as well as spatial inconsistency of crop (which increased). A distance incline is present in both areas, for couple of year, couple of *V. faba* variety, as well as couple of related bee-hive administration approaches. The average value is calculated is a 16 percent increase in yield, with 91 percent of that occurring within 766 metres of the hives. Providing hives are lucrative with a extensive variety of practical yield prices (dollars / tonne) as well as pollination costs, according to our economic review (dollars per hive) [6].

Methodology

Research Design:

It created a cause-as well as-effect model (Fig. 3 for an instance involving apple plants) to see in what way honeybee gathering administration influenced blossom inspection volume, that in turn influenced fruit set as well as quality. Three variables were used to measure the impact of gathering administration: a) honey bee gathering thickness in a 201-meter range, b) separation between the nearest gathering and a local tree, and c) gathering excellence (conservative vs. superiority). The quantity of florae on a tree is used as a metric for determining its attractiveness. It also looked at the quantity of tree variations in all plot to see if cross-pollination was a possibility. The quantity of seeds, which is proportional to pollination eminence; honey content; as well as d) distinct fruit bulk, that is a key aspect in the monetary assessment of the harvest as well as is also strongly associated to fruit diameter, distance, and extent.

Data Collection:

It tested causal hypotheses between variables using technique commonly utilised in ecosystem. Multi-level data were not taken into consideration in traditional route analysis. However, since many plants fitted to the identical farm, our model included multi-level data, and the farm feature had to be treated. To address the issue, researchers created a new path analysis technique named "Generalized multilevel path analysis," which takes into account the data's hierarchical structure. To summarise, the initial examination found entirely k likely "broken path," or variables that is not explicitly connected then consequently predictable to be statistically autonomous. If A cause B as well as B cause C, for example, the absent direction (k) is A's direct impact on C. Since correcting for the influence, the test involves calculating the likelihood (Pi) that A does not affect directly on C. The cumulative likelihood of all incomplete direction (k) of the track drawing is consequently determined to verify any proposed path analysis:

$$C = -2 \sum_{i=1}^K \ln(P_i)$$

Data Analysis:

C statistics has an approximated distribution. If the P value is less than 0.05, the path model is dismissed, indicating that considering the regulated indirect influence of A on C by B, there is also a straight consequence of A upon C. It used the Variance Inflation Factor to search in order to hypothesis the prototypical using Shipley's method. Then, using Scores, all variables were standardised. Finally, it used linear mixed-effect models to create the models aligned with each of the paths suggested in Fig. 3. It followed the method outlined.

Each test's residuals were tested for normal distribution. The Piecewise structural equation demonstrating kit performs comprehensive multilevel track investigates as well as calculates the C statistic, its scale, as well as the straight consequence for entirely variables, as well as providing a statistical importance and estimation of the C statistic (approximations of variables linked by an arrow as well as the p-value). Lastly, it looked at the impact of gathering efficiency upon apple as well as pear farmer incomes. The benefit discrepancies amid woods through tall- vs. traditional gatherings for the couple of fruit crop is associated using a non-parametric Wilcoxon-test to achieve this aim. When the Wilcoxon-test was important, it predicted ΔP .

Results And Discussion

It counted 1060 pollinator visit to pear as well as apple plants during the 260 pollinator censuses. *A. mellifera* made all of the visits, with the exception of. During nearly a couple of month study during apple as well as pear blossoming, it did not see uninhabited bee in the woods. As a result, *A. mellifera* is the sole source of pollination for these fruit crops. With a mean CI95 percent of .81, .16 as well as .25, .07 visit for 108 flowers for pear as well as apple flowers, respectively ($W = 260$, $P = .002$; Fig. 2), the inspection proportion is meaningfully advanced in apple flowers than in pear flower. The median is represented by the silver streak in the case, and the mean is represented by the white point. The third as well as first quartiles, respectively, are the lowest and highest confines of each case.




Fig. 2: Pear and Apple Trees Have A High Number of Visitors.

Our data is adequately explained by the generalised multilevel route model for apple trees (Fisher's C is equal to 39.99, k is equal to 30, P is greater than 0.04). The path diagram it evaluated is shown in Fig. 3, and their impact of colony efficiency upon honey bee inspection rate (estimate is equal to 0.75, SE is equal to 0.15, P is equal to 0.002) was the most notable finding. The number of fruits (estimate is equal to 0.24, SE is equal to 0.12, P is

equal to 0.05), the number of seeds (estimate is equal to 0.5, SE is equal to 0.15, P is equal to 0.004), and the pulp sugar concentration (estimate is equal to 0.45, SE is equal to 0.14, P is equal to 0.004) were all significantly affected by visitation rate. While bee visitation increased fruit quantity and consistency, (Estimate is equal to 0.78, SE is equal to 0.18, p 0.002) as well as individual fruit weight (Estimate is equal to 0.84, SE is equal to 0.17, p 0.002).

The magnitude of an impact is represented by the thickness of an arrow, which is also given. the arrow and sign of the figures indicate whether the *p 0. 04.. p 0.2. ***p 0.002, **p 0.02, *p 0. 04.. p 0.2. flow: measured number of florae/trees; col qual: colony quality; var: quantity of varieties/parcel. Farmers who rented high-quality colonies made significantly more money than farmers who rented traditional colony (W is equal to 65, P is equal to 0.044). Benefit aimed at farm using good colony is 16.9, 539.9, 10.9, 194.9 US\$/ha, while it is just 9.9, 259.9, 6.9, 86.9 US\$/ha for farms using traditional colonies. Agriculturalists who utilized superior colony received a marginal value (i.e. ΔP) of 7, 280 US \$/ha (Fig. 4).




Fig. 3: For Apple Trees, It Used A Generalised Multilevel Path Analysis.

(a) Traditional, the 108 blossom/min. (b) honey bee (108 bees). (c) apple sugar concentration (brixsi unit) in comparison to honey bee visitation intensity (108 bees/blossom/min). (d) apple as a function of fruit load/tree. (e) the weight of each apple (f) amount of fruit produced/tree (g) the impact (in US dollars) traditional represents the median, as well as the black point represents the mean, respectively, the black line represents the linear model's prediction, while the spotted outlines represent the 94.99 percent confidence pauses.

Fisher's P is greater than 0.049, k is equal to 29.9, C is equal to 39.85, the proposed generalised pear tree of the suits the results well. The number of fruits per tree, on the other hand, was only compared to the quantity of florae (P is equal to 0.045, SE is equal to 0.23, estimate is equal to 0.50). Visitation patterns were unaffected by colony management, which included 201-meter efficiency. As a result, it found no evidence of a connection between colony efficiency and farmer benefit (W is equal to 85, P is equal to 0.55). The suggested generalised (P is greater than 0.05, k is equal to 35, C is equal to 46.55) suits the results well (P is greater than 0.049, k is equal to 35, C is equal to 46.55). Individual fruit weight was significantly influenced by colony content (P is equal to 0.036, SE is equal to 0.20, estimate is equal to 0.479) but (P is equal to 0.030, SE is equal to 0.15, estimate is equal to 0.40). In this variety, major impact on a agriculturalist's benefit, as it did in previous result.




Fig. 4. Visitation, Fruit Production, And Farmer Benefit in Apple Are Affected by Colony Qualities, Inspection Volume, Besides Fruit Load.

Conclusion

Our research found that simple improvements leased honey bee colony would increase visitation rates, increasing and thus increasing agriculturalists' incomes outpacing availability of honey bee colonies, our findings indicate practises intended to enhancing cluster efficiency could be able to help address this possible colony number shortfall. *A. mellifera* is the sole pollinator. With the reported reduction of bees in world, condition like this could sadly spread. Despite widespread interest in pollinators' position in nutriment manufacture, their effect upon agriculturalists' benefit, that decides their subsistence as well as land-utilization choices, remains unknown. While normal pollinator assistances were commonly presumed, even within the same region, here was the latent for significant longitudinal distinction amongst yield types besides variations, as well as amongst pollinator administration policies.

It investigated efficiency of honey bee colony utilized for pollination service, that includes pathogen control and non-natural winter feeding, affects flowers visitation, farmer benefit, and fruit production (Patagonia). Flower visiting rates were 130.5 percent higher in high-quality apple colonies than in traditional colonies resulting in a 70.5% rise in farm income. Colony consistency only influenced fruit weight, varieties a rise in farmer income to the same degree as the apple. It found no, in comparison to other reports, demonstrating the vulnerability of this conventionally intensified crop production mechanism. The feasibility of existing pollination procedures must be tested as a first step can be influenced by improving pollination through successful management.

Pollination is important not only for total yield, but also for primary controlled, little is known about honey bees' contribution to farmer incomes. Understanding the importance of essential for developing management strategies that improve pollination while reducing production variability and farmer earnings. However, to determine impacts of pollination service, it is essential to relate pollination activities to a farmer's benefit. The majority of research to date have concentrated on the impact of various pollinator control systems on yield quantity and efficiency, with only a handful addressing the economic implications.

References

1. M. A. Aizen and L. D. Harder, "The Global Stock of Domesticated Honey Bees Is Growing Slower Than Agricultural Demand for Pollination," *Curr. Biol.*, 2009, doi: 10.1016/j.cub.2009.03.071.
2. M. A. Aizen, L. A. Garibaldi, S. A. Cunningham, and A. M. Klein, "Long-Term Global Trends in Crop Yield and Production Reveal No Current Pollination Shortage but Increasing Pollinator Dependency," *Curr. Biol.*, 2008, doi: 10.1016/j.cub.2008.08.066.
3. T. D. Breeze et al., "Agricultural policies exacerbate honeybee pollination service supply-demand mismatches across Europe," *PLoS One*, 2014, doi: 10.1371/journal.pone.0082996.
4. C. Brittain, C. Kremen, A. Garber, and A. M. Klein, "Pollination and plant resources change the nutritional quality of almonds for human health," *PLoS One*, 2014, doi: 10.1371/journal.pone.0090082.
5. M. Buccheri and C. Di Vaio, "Relationship Among Seed Number, Quality, and Calcium Content in Apple Fruits," *J. Plant Nutr.*, 2005, doi: 10.1081/pln-200026409.
6. S. A. Cunningham and D. Le Feuvre, "Significant yield benefits from honeybee pollination of faba bean (*Vicia faba*) assessed at field scale," *F. Crop. Res.*, 2013, doi: 10.1016/j.fcr.2013.05.019.
7. S. A. Cunningham, A. Fournier, M. J. Neave, and D. Le Feuvre, "Improving spatial arrangement of honeybee colonies to avoid pollination shortfall and depressed fruit set," *J. Appl. Ecol.*, 2016, doi: 10.1111/1365-2664.12573.
8. N. Deguines, C. Jono, M. Baude, M. Henry, R. Julliard, and C. Fontaine, "Large-scale trade-off between agricultural intensification and crop pollination services," *Front. Ecol. Environ.*, 2014, doi: 10.1890/130054.
9. E. J. Eilers, C. Kremen, S. S. Greenleaf, A. K. Garber, and A. M. Klein, "Contribution of pollinator-mediated crops to nutrients in the human food supply," *PLoS One*, 2011, doi: 10.1371/journal.pone.0021363.
10. R. Földesi et al., "Relationships between wild bees, hoverflies and pollination success in apple orchards with different landscape contexts," *Agric. For. Entomol.*, 2016, doi: 10.1111/afe.12135.